

The motion equation for massive fermions and bosons without Higgs

Gunn Quznetsov

454016, Chelyabinsk-16, yD.BET. Φ N 949892, Russia

email: lak@cgu.chel.su, gunn@mail.ru, quznets@yahoo.com

Abstract

The motion equation with nonzero mass, invariant for weak isospin transformation, can be obtained from the Dirac equation by the adding of the Clifford pentad fifth element. The motion equation of the SU(2) Yang-Mills field components is similar to the Klein-Gordon equation with the nonzero mass.

PACS: 13.10, 12.10, 12.15, 14.70.Fm, 14.80.B, 14.80.C

1 Introduction

I consider the motion equations for fermions with nonzero masses without Higgs with all five elements of Clifford's pentad. And i reform the motion equation for components W-field as equation, similar to the nonzero Klein-Gordon equation.

Notations:

$$1_2 \stackrel{Def}{=} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, 1_4 \stackrel{Def}{=} \begin{bmatrix} 1_2 & 0_2 \\ 0_2 & 1_2 \end{bmatrix},$$

$$0_2 \stackrel{Def}{=} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, 0_4 \stackrel{Def}{=} \begin{bmatrix} 0_2 & 0_2 \\ 0_2 & 0_2 \end{bmatrix}$$

$$\gamma^{[5]} \stackrel{Def}{=} \begin{bmatrix} 1_2 & 0_2 \\ 0_2 & -1_2 \end{bmatrix}, \beta^{[0]} \stackrel{Def}{=} -1_4, 1_8 \stackrel{Def}{=} \begin{bmatrix} 1_4 & 0_4 \\ 0_4 & 1_4 \end{bmatrix},$$

the Pauli matrices:

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix};$$

if ϑ is the 4×4 matrix then

$$\underline{\vartheta} \stackrel{Def}{=} \begin{bmatrix} \vartheta & 0_4 \\ 0_4 & \vartheta \end{bmatrix}.$$

A set $\tilde{C}$ of complex $n \times n$ matrices is denoted as *Clifford's set* [3] if the following conditions are fulfilled:

if $\alpha_k \in \tilde{C}$ and $\alpha_r \in \tilde{C}$ then $\alpha_k \alpha_r + \alpha_r \alpha_k = 2\delta_{k,r}$;

if $\alpha_k \alpha_r + \alpha_r \alpha_k = 2\delta_{k,r}$ for all elements α_r of $\tilde{C}$ then $\alpha_k \in \tilde{C}$.

If $n = 4$ then Clifford's set either contains 3 matrices (*Clifford's triplet*) or contains 5 matrices (*Clifford's pentad*).

For example: *Clifford pentad* β :

$$\beta^{[1]} \stackrel{Def}{=} \begin{bmatrix} \sigma_1 & 0_2 \\ 0_2 & -\sigma_1 \end{bmatrix}, \beta^{[2]} \stackrel{Def}{=} \begin{bmatrix} \sigma_2 & 0_2 \\ 0_2 & -\sigma_2 \end{bmatrix}, \beta^{[3]} \stackrel{Def}{=} \begin{bmatrix} \sigma_3 & 0_2 \\ 0_2 & -\sigma_3 \end{bmatrix}, \quad (1)$$

$$\gamma^{[0]} \stackrel{Def}{=} \begin{bmatrix} 0_2 & 1_2 \\ 1_2 & 0_2 \end{bmatrix}, \quad (2)$$

$$\beta^{[4]} \stackrel{Def}{=} i \begin{bmatrix} 0_2 & 1_2 \\ -1_2 & 0_2 \end{bmatrix}. \quad (3)$$

2 Fermions

Let the motion equation (Dirac equation) for the free fermion be in the following form:

$$\left(\underline{\beta}^{[0]} i \partial_0 + \underline{\beta}^{[1]} i \partial_1 + \underline{\beta}^{[2]} i \partial_2 + \underline{\beta}^{[3]} i \partial_3 + \underline{\gamma}^{[0]} m \right) \psi = 0. \quad (4)$$

The motion equation for two independent fermions with masses m_1 and m_2 (a two-fermion state) is of the following:

$$\left(\underline{\beta}^{[0]} i \partial_0 + \underline{\beta}^{[1]} i \partial_1 + \underline{\beta}^{[2]} i \partial_2 + \underline{\beta}^{[3]} i \partial_3 + \underline{\gamma}^{[0]} m_{1,2} \right) \Psi = 0$$

with

$$m_{1,2} \stackrel{Def}{=} \begin{bmatrix} m_1 \cdot 1_4 & 0_4 \\ 0_4 & m_2 \cdot 1_4 \end{bmatrix}.$$

This equation is not invariant for the SU(2) isospin transformation

$$U \stackrel{Def}{=} \begin{bmatrix} (a + ib) 1_2 & 0_2 & (c + iq) 1_2 & 0_2 \\ 0_2 & 1_2 & 0_2 & 0_2 \\ (-c + iq) 1_2 & 0_2 & (a - ib) 1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 1_2 \end{bmatrix}$$

with

$$a^2 + b^2 + c^2 + q^2 = 1$$

because it holds four, only, elements of the Clifford pentad.

If $\partial_\mu U = U \partial_\mu$ then the equation with all five elements

$$\left(\underline{\beta^{[0]}} i \partial_0 + \underline{\beta^{[1]}} i \partial_1 + \underline{\beta^{[2]}} i \partial_2 + \underline{\beta^{[3]}} i \partial_3 + \underline{\gamma^{[0]}} m_5 + \underline{\beta^{[4]}} m_4 \right) \Psi = 0$$

is invariant for the following transformation:

$$\begin{aligned} \Psi &\rightarrow \Psi' = U \Psi; \\ m_5 &\rightarrow m'_5 = a m_5 (\ell_o + \ell_o) - \sqrt{1 - a^2} m_4 (\ell_o - \ell_o); \\ m_4 &\rightarrow m'_4 = a m_4 (\ell_o + \ell_o) + \sqrt{1 - a^2} m_5 (\ell_o - \ell_o) \end{aligned}$$

with

$$\begin{aligned} \ell_o &\stackrel{Def}{=} \frac{1}{2\sqrt{1-a^2}} \begin{bmatrix} \left(b + \sqrt{1-a^2} \right) 1_4 & (q - ic) 1_4 \\ (q + ic) 1_4 & \left(\sqrt{1-a^2} - b \right) 1_4 \end{bmatrix}, \\ \ell_* &\stackrel{Def}{=} \frac{1}{2\sqrt{1-a^2}} \begin{bmatrix} \left(\sqrt{1-a^2} - b \right) 1_4 & (-q + ic) 1_4 \\ (-q - ic) 1_4 & \left(b + \sqrt{1-a^2} \right) 1_4 \end{bmatrix}. \end{aligned}$$

In this transformation the value of $m_5^2 + m_4^2$ is invariant.

Hence we have got the two-fermion motion equation, invariant for global weak isospin transformation, with nonzero nonHiggs mass.

If g is a real number, $W_\mu^0, W_\mu^1, W_\mu^2$, are real scalar functions on the 3+1 space-time and

$$W_\mu \stackrel{Def}{=} \begin{bmatrix} W_\mu^0 1_2 & 0_2 & (W_\mu^1 - i W_\mu^2) 1_2 & 0_2 \\ 0_2 & 1_2 & 0_2 & 0_2 \\ (W_\mu^1 + i W_\mu^2) 1_2 & 0_2 & -W_\mu^0 1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 1_2 \end{bmatrix}$$

then the equation

$$\left(\sum_{\mu=0}^3 \underline{\beta^{[\mu]}} i \left(\partial_\mu - i \frac{1}{2} g W_\mu \right) + \underline{\gamma^{[0]}} m_5 + \underline{\beta^{[4]}} m_4 \right) \Psi = 0$$

is invariant for the local weak isospin transformation:

$$\begin{aligned} \Psi &\rightarrow \Psi' = U \Psi; \\ m_5 &\rightarrow m'_5 = a m_5 (\ell_o + \ell_*) - \sqrt{1 - a^2} m_4 (\ell_o - \ell_*); \\ m_4 &\rightarrow m'_4 = a m_4 (\ell_o + \ell_*) + \sqrt{1 - a^2} m_5 (\ell_o - \ell_*), \\ W_\mu &\rightarrow W'_\mu = U W_\mu U^\dagger - \frac{2i}{g} (\partial_\mu U) U^\dagger. \end{aligned}$$

3 Bosons

The motion equation of the Yang-Mills SU(2) field in the space without matter (for instance [5] or [4]) has got the following form:

$$\partial^\nu \mathbf{W}_{\mu\nu} = -g \mathbf{W}^\nu \times \mathbf{W}_{\mu\nu}$$

with:

$$\mathbf{W}_{\mu\nu} = \partial_\mu \mathbf{W}_\nu - \partial_\nu \mathbf{W}_\mu + g \mathbf{W}_\mu \times \mathbf{W}_\nu$$

and

$$\mathbf{W}_\mu = \begin{bmatrix} W_\mu^0, \\ W_\mu^1, \\ W_\mu^2, \end{bmatrix}.$$

Hence the motion equation for W_μ^0 , is the following:

$$\begin{aligned} \partial^\nu \partial_\nu W_\mu^0 = & g^2 (W^{2,\nu} W_\nu^2 + W^{1,\nu} W_\nu^1) W_\mu^0 - \\ & - g^2 (W^{1,\nu} W_\mu^1 + W^{2,\nu} W_\mu^2) W_\nu^0 + \\ & + g \partial^\nu (W_\mu^1 W_\nu^2 - W_\mu^2 W_\nu^1) + \\ & + g (W^{1,\nu} \partial_\mu W_\nu^2 - W^{1,\nu} \partial_\nu W_\mu^2 - W^{2,\nu} \partial_\mu W_\nu^1 + W^{2,\nu} \partial_\nu W_\mu^1) + \\ & + \partial^\nu \partial_\mu W_\nu^0. \end{aligned} \quad (5)$$

W_μ^1 , and W_μ^2 , satisfy to similar equations.

This equation can be reformed as the following:

$$\begin{aligned} \partial^\nu \partial_\nu W_\mu^0 = & [g^2 (W^{2,\nu} W_\nu^2 + W^{1,\nu} W_\nu^1 + W^{0,\nu} W_\nu^0)] \cdot W_\mu^0 - \\ & - g^2 (W^{1,\nu} W_\mu^1 + W^{2,\nu} W_\mu^2 + W^{0,\nu} W_\mu^0) W_\nu^0 + \\ & + g \partial^\nu (W_\mu^1 W_\nu^2 - W_\mu^2 W_\nu^1) + \\ & + g (W^{1,\nu} \partial_\mu W_\nu^2 - W^{1,\nu} \partial_\nu W_\mu^2 - W^{2,\nu} \partial_\mu W_\nu^1 + W^{2,\nu} \partial_\nu W_\mu^1) + \\ & + \partial^\nu \partial_\mu W_\nu^0. \end{aligned}$$

This equation looks like to the Klein-Gordon equation of field W_μ^0 , with mass

$$g [- (W^{2,\nu} W_\nu^2 + W^{1,\nu} W_\nu^1 + W^{0,\nu} W_\nu^0)]^{\frac{1}{2}}. \quad (6)$$

and with the additional terms of the W_μ^0 , interactions with others components of $\mathbf{W}$.

"Mass" (6) is invariant for the following transformations:

$$\begin{cases} W_r^{k,\prime} = W_r^k \cos \lambda - W_s^k \sin \lambda, \\ W_s^{k,\prime} = W_r^k \sin \lambda + W_s^k \cos \lambda; \end{cases} \quad \left| \begin{cases} W_0^{k,\prime} = W_0^k \cosh \lambda - W_s^k \sinh \lambda, \\ W_s^{k,\prime} = W_s^k \cosh \lambda - W_0^k \sinh \lambda \end{cases} \right|$$

with a real number λ , and $r \in \{1, 2, 3\}$, and $s \in \{1, 2, 3\}$,
and (6) is invariant for a global weak isospin transformation U :

$$W_\nu \rightarrow W'_\nu = UW_\nu U^\dagger$$

but is not invariant for a local transformation U :

$$W_\nu \rightarrow W'_\nu = UW_\nu U^\dagger - \frac{2i}{g} (\partial_\nu U) U^\dagger.$$

Equation (5) can be simplified as follows:

$$\begin{aligned} \sum_\nu g_{\nu,\nu} \partial^\nu \partial_\nu W_\mu^0 = & \left[g^2 \sum_{\nu \neq \mu} g_{\nu,\nu} \left((W_\nu^2)^2 + (W_\nu^1)^2 \right) \right] \cdot W_\mu^0 - \\ & - g^2 \sum_{\nu \neq \mu} g_{\nu,\nu} (W_\mu^{1,\nu} W_\mu^1 + W_\mu^{2,\nu} W_\mu^2) W_\nu^0 - \\ & + g \sum_\nu g_{\nu,\nu} \partial^\nu (W_\mu^1 W_\nu^2 - W_\mu^2 W_\nu^1) + \\ & + g \sum_\nu g_{\nu,\nu} (W_\mu^{1,\nu} \partial_\mu W_\nu^2 - W_\mu^{2,\nu} \partial_\mu W_\nu^1 + W_\mu^{2,\nu} \partial_\nu W_\mu^1) + \\ & + \partial_\mu \sum_\nu g_{\nu,\nu} \partial^\nu W_\nu^0. \end{aligned}$$

(here no of summation over indexes ν ; the summation is expressed by $\sum$)
with $g_{0,0} = 1$, $g_{1,1} = g_{2,2} = g_{3,3} = -1$.

In this equation the form

$$g \left[- \sum_{\nu \neq \mu} g_{\nu,\nu} \left((W_\nu^2)^2 + (W_\nu^1)^2 \right) \right]^{\frac{1}{2}}$$

varies in space, but it does not contain W_μ^0 and locally acts as mass - i.e. it does not allow to particles of this field to behave as a massless ones.

Hence if Higgs exist then mass, calculated with Higgs, should be corrected with this "mass". If Higgs do not exist then this "mass" turns out to be the only reason, giving a W -boson nonzero mass in measurements [2].

Therefore the W fields should behave as the massive fields with Higgs or without Higgs.

4 Conclusion

Therefore Higgs are not necessary for the nonzero leptons' and bosons' masses.

References

- [1] for instance, Gordon Kane, *Modern Elementary Particle Physics*.
Addison-Wesley Publishing Company, Inc., (1987)
- [2] for instance, Klaus Moenig *Preprint* hep-ph/0303023
- [3] for instance, E. Madelung, *Die Mathematischen Hilfsmittel des Physikers*.
(Springer Verlag, 1957) p.29

- [4] Lewis H. Ryder, *Quantum Field Theory*, Mir, Moscow (1987), p.133, (3.136)
- [5] M. V. Sadoyski, *Lectures on the Quantum Fields Theory* (Institute of Electrophysics, UrO RAS, 2000), p.33 (2.126)